



# MYCORRHIZA NEWS

The Quarterly Newsletter of Mycorrhiza network

Volume 25 • Issue 4 • January 2014



## About TERI

TERI (The Energy and Resources Institute) is a dynamic and flexible organization with a global vision and a local focus. TERI's focus is on research, in the fields of energy, environment, and sustainable development, and on documentation and information dissemination. The genesis of these activities lie in TERI's firm belief that the efficient utilization of energy, sustainable use of natural resources, large-scale adoption of renewable energy technologies, and reduction of all forms of waste would move the process of development towards the goal of sustainability.

## TERI's Mycorrhiza Network

TERI's Mycorrhiza Network is primarily responsible for establishing the MIC (Mycorrhiza Information Centre), the CMCC (Centre for Mycorrhiza Culture Collection), and publishing Mycorrhiza News. The Network helps scientists carry out research in mycorrhiza and promotes communication among mycorrhiza scientists.

## *Mycorrhiza News*

The Mycorrhiza News provides a forum for dissemination of scientific information on mycorrhiza research and activities; publishes state-of-the-art papers from eminent scientists; notes on important breakthroughs; brief accounts of new approaches and techniques; publishes papers compiled from its RIZA database; provides information on forthcoming events on mycorrhiza and related subjects; lists important research references published during the quarter; and highlights the activities of the CMCC.



# CONTENTS

## RESEARCH FINDING PAPERS

- Arbuscular Mycorrhizal Colonization in *Aquilaria malaccensis* Lamk. and in its Surrounding Herbaceous Community 2
- Selection of Efficient VAM Species for Marigold Cultivation and its Effect on Growth, Yield, and Quality Parameters of two Marigold Varieties 8

## CMCC ARTICLE

- Morphotaxonomy of Arbuscular Mycorrhizal Fungi *Acaulospora foveata* 14

- RECENT REFERENCES 17

- FORTHCOMING EVENTS 20

## RESEARCH FINDING PAPERS

### Arbuscular Mycorrhizal Colonization in *Aquilaria malaccensis* Lamk. and in its Surrounding Herbaceous Community

Atithi Debnath<sup>1</sup>, Ajay Krishna Saha<sup>2</sup>, and Panna Das<sup>\*</sup>

#### Introduction

An increasing demand for herbal products has endangered many traditionally used and pharmaceutically important plant species and their habitats (Fuchs and Haselwandter, 2008). *Aquilaria malaccensis* Lamk. popularly known as agarwood belongs to the family Thymelaeaceae and class Magnoliopsida. It is one of the world's most expensive essential oil yielding tree species. Most of the *Aquilaria* species are found in south-east Asia (Donovan and Puri, 2004). *A. malaccensis* and *A. khasiana* are reported to grow naturally in the plains and in the foothills of north-east India (Hajra, 2000). The oleoresin contained in its wood is widely used in the perfume industry and in medicines (Sumadiwangsa, 1997). Studies have revealed that agarwood has remarkable anticancer activity (Gunasekera *et al.* 1981). A rise in demand for agarwood has resulted in irrational cutting of the tree trunk for extraction of the chemical. Hence, the tree has become endangered now.

Arbuscular Mycorrhizal (AM) fungi are ubiquitous symbionts existing in any ecosystem and colonizing in over two-thirds of the vascular plant species (Koide and Mosse, 2004). AM fungi are naturally occurring soil fungi belonging to a recently new ascribed phylum, the Glomeromycota. This phylum is presumed to have originated at least 460

million years ago (Schüßler *et al.* 2001). AM fungi stimulate uptake of plant nutrients, such as P, Zn, Cu, and Fe in deficient soils. Also, mycorrhizal hyphae can significantly improve N, P, and K uptake (Chen and Zhao, 2009).

The objective of this study was to analyze AM fungal colonization and diversity of *A. malaccensis* and its surrounding herbaceous community growing in Tripura, north-east India.

#### Materials and Methods

##### Sample Collection

Roots from five plants species, viz., *A. malaccensis*, *Clerodendrum viscosum* Vent., *Phaulopsis dorsiflora* (Retz.) Sant., *Eupatorium odoratum* L., and *Mimosa pudica* L. were collected from Syamalibazar area, Agartala, Tripura. Sample collection was carried out from August 2012 to December 2012. Root samples were collected using a trowel to dig a constant maximum depth of 10cm. Most of the fine roots of the plants were located in the upper 10cm of the soil profile. Roots were traced to their origin to ensure that they were from the desired plants.

The rhizospheric soil, at depths of 10cm, surrounding the roots of the *A. malaccensis* tree was collected from eight different points. The rhizospheric soils from the roots of the five herbaceous plants were also collected. The combined samples of

<sup>1</sup>Microbiology Laboratory, Department of Botany, Tripura University, Suryamaninagar-799 022, Tripura, India

<sup>2</sup>Mycology and Plant Pathology Laboratory, Department of Botany, Tripura University, Suryamaninagar-799 022, Tripura, India

<sup>\*</sup>Corresponding author. E-mail panna11d@yahoo.com

approximately 500 g soil of *A. malaccensis* and all the soil samples of herbaceous plants mixed, laced in polythene bags, labelled, and were transported to the laboratory for further analysis.

#### Preparation of Roots and Assessment of AMF

Collected roots were thoroughly washed with tap water several times and cut into uniform pieces approximately 1 cm long. Then the roots were cleaned with 10 per cent NaOH. The cleaned roots were again washed with tap water and bleached in two drops of alkaline H<sub>2</sub>O<sub>2</sub> before acidification for 2–3 minutes. After acidification, the roots were again washed and stained with black Faber Castell stamp pad ink (Das and Kayang, 2008). These root segments were then mounted on slide and examined for mycorrhizal structures under compound microscope (Olympus 108210).

#### Spore Analysis

The spores were extracted by modified wet sieving and decanting method (Muthukumar, Senthikumar, and Rajangam *et al.* 2006). The isolated spores were transferred using a wet needle to polyvinyl alcohol-lactoglycerol (PVLG) under a dissecting microscope (Koske and Tessier, 1983) for identification. The complete and broken spores were examined using a compound microscope (Olympus 108210). The taxonomic identification of spores was based on spore size, colour, ornamentation, and wall characteristics by matching original descriptions.

#### Soil Analysis

Soil moisture was determined by drying 10 g of fresh soil at 60 °C for 24 hours in a hot-air oven. For pH and electrical conductivity, 10 g of soil was dissolved in 50 ml of distilled water and stirred for 20 minutes. This solution was kept overnight and then the soil pH and electrical conductivity was measured using a digital meter. The organic carbon, available phosphorus, potassium, and calcium contents of the soil were determined using the Jackson (1978) method. The available nitrogen in the soil was estimated by Black (1982) method.

#### Data Analysis

Standard error of means and Tukey test was conducted to separate the means in Statistica 9.0.

Diversity indices were calculated (Hammer, Harper, Ryan *et al.* 2001).

## Results and Discussion

#### Soil Properties

Table 1 depicts the properties of soils collected from the rhizosphere of *Aquilaria malaccensis* and herbaceous plants. Moisture content of soil supporting *A. malaccensis* was higher than the soils harbouring herbaceous plants. The soil pH was found to be acidic while the electrical conductivity was much higher in *A. malaccensis* than herbaceous plants' soil. The other properties of soil, such as organic carbon per cent, available nitrogen, available phosphorus, and available potassium were higher in soils from *A. malaccensis* than the soils of herbaceous plants. The available calcium was higher in soils from herbaceous plants than soils supporting *A. malaccensis*. Nutrient concentration, pH, and soil humidity level can influence fungal distribution, root colonization, and mycorrhizal efficiency (Koide and Li, 1990). In general, soils having pH 6.0–6.3 support greater number of AM propagules than soils having pH 5.3–5.7 (Tabin *et al.* 2009).

#### Root Colonization by AM fungi

All the root samples collected from the tree and the surrounding herbaceous plants exhibited the presence of AM colonization showing arbuscules, vesicles, and hyphae. The mycorrhizal structural colonization in the roots of *Aquilaria malaccensis* and its surrounding herbaceous plants is presented in Table 2. The percentage of arbuscule is highest in *A. malaccensis* and lowest in *Eupatorium odoratum*. The percentage of vesicle was highest in *Mimosa pudica* and lowest in *Clerodendrum viscosum*. The percentage of hyphae was highest in *A. malaccensis* and lowest in *Phaulopsis dorsiflora* (Fig. 1). AM fungal colonization in *A. malaccensis* and its surrounding herbaceous community forms the first report. Similar results were reported in *A. malaccensis* by Tabin *et al.* (2009). Arbuscular, vesicular, and hyphal colonization were found in all the species. The percentage of mycorrhizal colonization showed a variation from 3.86 per cent to 87.88 per cent.

**Table 1:** Properties of soils collected from the rhizosphere of *Aquilaria malaccensis* and herbaceous plants

Soil Samples	Moisture Content (%)	pH	Electrical Conductivity (mS cm <sup>-1</sup> )	Organic Carbon (%)	Available Nitrogen (Kg/ha)	Available Phosphorus (Kg/ha)	Available Potassium (Kg/ha)	Available Calcium (Kg/ha)
<i>Aquilaria malaccensis</i>	11.5	4.86	119	0.75	13.84	0.35	11.37	10.73
Herbaceous plants	7.32	5.84	49	0.59	8.44	0.23	8.55	14.93

**Table 2:** AM fungal colonization (%) in the roots of *Aquilaria malaccensis* and its surrounding herbaceous plants

Name of Plants	% RLA <sup>a</sup>	% RLV <sup>a</sup>	% RLH <sup>a</sup>
<i>Aquilaria malaccensis</i>	20.72±3.77 <sup>a</sup>	11.50±2.85 <sup>a</sup>	87.88±2.85 <sup>a</sup>
<i>Clerodendrum viscosum</i>	11.15±2.92 <sup>b</sup>	8.71±1.79 <sup>a</sup>	69.28±4.54 <sup>b</sup>
<i>Phaulopsis dorsiflora</i>	4.56±1.45 <sup>b</sup>	8.89±2.98 <sup>a</sup>	50.86±3.30 <sup>c</sup>
<i>Eupatorium odoratum</i>	3.86±1.21 <sup>b</sup>	17.17±2.82 <sup>a</sup>	61.56±3.70 <sup>b</sup>
<i>Mimosa pudica</i>	4.68±1.56 <sup>b</sup>	31.29±3.60 <sup>b</sup>	55.16±3.41 <sup>b</sup>

Tukey test showing that different alphabetical letters are significantly different ( $p < 0.05$ )


%RLA, %RLV, and %RLH are percent root length with arbuscules, vesicles, and hyphae, respectively.

### AM Fungal Species


Figure 2 depicts the AM fungal spore density in 25 g/ soil sample. Seven different AM fungal species were found in both the soil samples. These were *Glomus*

*glomerulatum*, *Claroideoglomus etunicatum*, *Glomus* sp 1, *Glomus* sp 2, *Glomus* sp 3, *Glomus* sp 4, and *Acaulospora* sp 1 (Plate 1). These species of AM fungi were collected and identified on the basis of their

**Fig. 1:** Comparison of AM fungal colonization (%) between *Aquilaria malaccensis* and its surrounding herbaceous plants




**Fig. 2:** Spore density/25g soil of *Aquilaria malaccensis* and herbaceous plants



**Plate 1:** AM fungal species (a) *Glomus glomerulatum* (x400) (b) *Claroideoglomus etunicatum* (x400) (c) *Glomus* sp 1 (x400) (d) *Glomus* sp 2 (x400) (e) *Glomus* sp 3 (x400) (f) *Glomus* sp 4 (x400) (g) *Acaulospora* sp 1 (x400)





morphological characteristics. Among these species, *Glomus* sp 2 was common in both the samples. The abundance and relative abundance (in %) of AM fungi was given in Table 3. In herbaceous, *Glomus* sp 1 showed the highest abundance whereas the lowest abundance was by *Acaulospora* sp 1. In *A. malaccensis* *Glomus* sp 2 showed the highest abundance and the lowest was shown by *Glomus glomerulatum*. In the present study, the AMF spore density for *A. malaccensis* and herbaceous plants was found to be 32 and 33 spores/ 25 g soil, respectively, and the number of spores ranged from 11 to 384 spores per 100g soil reported in forest ecosystems of temperate regions (Read *et al.* 1976). Prasad K (1998) also reported a range of 5–370 spores/100 g of dried soil in India. The spore density was low due to the production of AMF spores in the rhizosphere vicinity of herbaceous plant species (Kruckelmann, 1975). Mohammad *et al.* (2003) observed that the spore production increased

with the soil pH and organic carbon. In the present study, the pH was low for both the soil samples which might have influenced the number of spores. Therefore, spore population is affected by a wide range of soil, climatic, fungal, and host factors (Anderson *et al.* 1983).

The diversity indices namely Dominance, Shannon, and Evenness showed no significant difference between the samples. The diversity index is presented in Table 4. Seven species of AMF were obtained from the soil samples of *A. malaccensis* and herbaceous plants, respectively, which were lower than the 8–20 species usually reported in arable lands (Jansa *et al.* 2002). Valsalakumar, Ray, and Potty (2007) also reported very low AMF diversities of one to three taxa for the 21 sampling locations from soil supporting *Phaseolus aureus* in Tamil Nadu and Karnataka, respectively. Out of the seven species, the genus *Glomus* was found to be predominant. Similar result was recorded by Tabin *et al.* 2009 in *Aquilaria*

**Table 3:** Relative abundance (in %) of AM fungal species extracted from the rhizosphere of herbaceous plant and *Aquilaria malaccensis*

AM fungi	Herbaceous plants		<i>Aquilaria malaccensis</i>	
	Abundance	Relative abundance (%)	Abundance	Relative abundance (%)
<i>Acaulospora</i> sp 1	2	6.90	-	-
<i>Claroideoglomus etunicatum</i>	4	13.79	-	-
<i>Glomus glomerulatum</i> sp. nov. Sieverding	-	-	4	14.81
<i>Glomus</i> sp 1	16	55.17	-	-
<i>Glomus</i> sp 2	7	24.14	11	40.74
<i>Glomus</i> sp 3	-	-	5	18.52
<i>Glomus</i> sp 4	-	-	7	25.93

**Table 4:** Diversity index of AM fungal species in herbaceous plants and *Aquilaria malaccensis*

Diversity Indices	Herbaceous Plants	<i>Aquilaria malaccensis</i>
Dominance_D	0.3864	0.2894
Shannon_H	1.129	1.311
Simpson_1-D	0.6136	0.7106
Evenness_e^H/S	0.773	0.9275

plant. Das and Kayang (2009) also reported the dominance of *Glomus* sp. in north-east region of India. Certain species of Glomales are adapted to acidic soils and generally, dominate the AMF community (Sieverding, 1991) or they may be dominant due to their high competitiveness and reproductive capability (Sieverding, 1991). The abundance of AMF species showed the same tendency. Four species of AM fungi were isolated from *A. malaccensis* which is less than nine species observed (Tabin, *et al.* 2009) in *A. malaccensis* from Arunachal Pradesh. Although, spore populations do not exactly reflect the AMF community colonizing the plant roots because there is a possible existence of some non-sporulating AMF species as well (Daniell *et al.*, 2001).

The present study showed that the multiplication of indigenous AM fungi associated with *A. malaccensis* and herbaceous plants may possibly help in the sustained growth of *A. malaccensis*.

## Acknowledgement

The first author is grateful to UGC for the financial assistance. The authors are thankful to Head, Department of Botany, Tripura University, for providing facilities.

## References

- Anderson R C, Liberta A E, Dickman L A, and Katz A J. 1983. **Spatial variation in vesicular arbuscular mycorrhiza spore density.** *Bulletin of Torrey Botanical Club* 110: 519–525
- Black C A. 1982. *Methods of soil analysis.* Pregmon Press, England
- Chen X H and Zhao B. 2009. **Arbuscular mycorrhizal fungi mediated uptake of nutrient elements by Chinese milk vetch (*Astragalus sinicus* L.) grown in lanthanum spiked soil.** *Biology and Fertility of Soils* 45: 675–678
- Daniell T J, Husband R, Fitter A H, and Young J P W. 2001. **Molecular diversity of arbuscular mycorrhizal fungi colonizing arable crops.** *FEMS Microbial Ecology* 36: 203–209
- Das P and Kayang H. 2008. **Stamp pad ink, an effective stain for observing arbuscular mycorrhizal structure in roots.** *World Journal of Agricultural Science* 4: 58–60.
- Das P and Kayang H. 2009. **Arbuscular mycorrhizal fungi association with *Blechnum orientale* Linn. in pine forest and anthropogenically disturbed areas of northeast India.** *Archives of Agronomy and Soil Sciences* 55: 623–632
- Donovan D G and Puri R K. 2004. **Learning from traditional knowledge of non-timber forest products: Penan Benalui and the autecology of *Aquilaria* in Indonesian Borneo.** *Ecology and Society* 9: 1–23.
- Fuchs B and Haselwandter K. 2008. **Arbuscular mycorrhiza of endangered plant species: Potential impacts on restoration strategies.** In: Varma A (ed) *Mycorrhiza: state of the art, genetics and molecular biology, eco-function, biotechnology, eco-physiology, structure and systematics.* Springer, Heidelberg
- Gunasekera S P, Kinghorn A D, Cordell G A, and Farnsworth N R. 1981. **Plant anticancer agents, XIX. Constituents of *Aquilaria malaccensis*.** *Journal of Natural Products* 44: 569–572
- Hajra P K. 2000. **A review of the genus *Aquilaria* Lamk. in India.** In: M. Ahmed, P. Gogoi & G.U. Ahmed (eds.) *Proceedings of Seminar on Scope & Dimension of Agar (*Aquilaria* spp.) Plantation in NE Region.* All Assam Traders & Agar Oil Manufacturers Association, Nagaon (Asom), pp1–5
- Hammer O, Harper D A T, and Ryan P D. 2001. **PAST: Paleontological statistics software package for education and data analysis.** *Palaeontologia Electronica* 4: 1–9
- International Culture Collection of (Vesicular) Arbuscular Mycorrhizal Fungi, West Virginal University, <http://www.invam.caf.wvu.edu>
- Jackson M L. 1978. *Soil Chemical Analysis.* Prentice Hall, New Delhi, India
- Jansa J, Mozafar A, Anken T, Ruh R, Sanders I R, and Frossard E. 2002. **Diversity and structure of AMF communities as affected by tillage in a temperate soil.** *Mycorrhiza* 12: 225–234
- Koide R and Li M. 1990. **On host regulation of the vesicular arbuscular mycorrhizal symbiosis.** *New Phytologist* 114: 59–74
- Koide R T and Mosse B. 2004. **A history of research on arbuscular mycorrhiza.** *Mycorrhiza* 14:145–163
- Koske R E and Tessier B. 1983. **A convenient, permanent slide mounting medium.** *Mycological Society of American Newsletter* 34: 59
- Kruckelmann H W. 1975. **Effect of fertilizers, soils, soil tillage and plant species on the frequency of Endogone chlamydosporos and mycorrhizal infections in arable soils.** In: P.E. Sanders, B. Mosse and P.B. Tinker (eds), *Endomycorrhizas.* London: Academic Press. pp469–484
- Leibniz-Rechenzentrum, Bayerische Akademie der Wissenschaften, <http://www.lrz.de/~schuessler/amphylo>
- Mohammad M J, Hamad S R, and Malkawi H I. 2003. **Population of arbuscular mycorrhizal fungi in semi-arid environment of Jordan as influenced by biotic and abiotic factors.** *Journal of Arid Environments* 53: 409–417
- Muthukumar T, Senthilkumar M, Rajangam M, and Udaiyan K. 2006. **Arbuscular mycorrhizal morphology**

**and dark septate fungal associations in medicinal and aromatic plants of Western Ghats, Southern India.**

*Mycorrhiza* 17: 11–24

Muthukumar T and Udaiyan K. 2002. **Seasonality of vesicular-arbuscular mycorrhizae in seages in a semi-arid tropical grassland.** *Acta Oecologia* 23: 337–347

Prasad K. 1998. **Survey of vesicular-arbuscular mycorrhizal (VAM) fungal colonization in some weed species.** *Indian Journal of Applied and Pure Biology* 13: 87–90

Read D J, Koucheiki H K, and Hodgson J. 1976. **Vesicular-arbuscular mycorrhiza in natural vegetation systems. I. The occurrence of infection.** *New Phytologist* 77: 641–643

Schüßler A, Schwarzott D, and Walker C. 2001. **A new fungal phylum, the Glomeromycota: Phylogeny and**

**evolution.** *Mycological Research* 105: 1413–1421

Sieverding E. 1991. **Vesicular-arbuscular mycorrhiza management in tropical agrosystems.** Eschborn, Germany.

Sumadiwangsa S. 1997. **Agarwood as a high-value commodity in East Kalimantan [Kayu gaharu komoditi elit di Kalimantan Timur].** *Duta Rimba* 20: 33–40

Tabin T, Arunachalam A, Shrivastava K, and Arunachalam K. 2009. **Effect of arbuscular mycorrhizal fungi on damping-off disease in *Aquilaria agallocha* Roxb. seedlings.** *Tropical Ecology* 50: 243–248

Valsalakumar N, Ray J G, and Potty V P. 2007. **Arbuscular mycorrhizal fungi associated with Green gram in Southern India.** *Argonomy Journal* 99: 1260–1264

# Selection of Efficient VAM Species for Marigold Cultivation and its Effect on Growth, Yield, and Quality Parameters of two Marigold Varieties

Vijetha Patil, Chaya P Patil\*, and B S Kulkarni

## Introduction

In the present paper, the authors have reported the effect of 12 different species of mycorrhizal inoculation on the germination and growth of marigold raised in pot with soil as the media. This experiment was conducted to know whether all the species of mycorrhizal inoculation would be beneficial for plant growth. Vesicular–arbuscular mycorrhizal (VAM) fungi are obligate parasites that form symbiotic relationships with plant roots. VAM fungi are associated with increased phosphorus uptake, growth, and with increased drought resistance due to a fungal-hyphae-extended nutrient uptake zone around roots of many species (Hayman, 1980). Different species of VAM have different growth promotional effects on particular plant species (Patil and Patil, 2008). Thus, when considering VAM inoculation, it is important to examine several different species of VAM for their effect on plant growth.

Marigold (*Tagetes erecta* L.), which occupies a prominent place in ornamental horticulture, is a commercially exploited flower crop originated from Mexico and belongs to the family Asteraceae. Free flowering habit, short duration to produce marketable flowers, wide spectrum of attractive colours, shape, size, and good keeping quality make marigold an attractive option for flower growers. Apart from its significance in ornamental horticulture, it is valued for its other properties also. The aromatic oil extracted from *Tagetes minuta*, commonly known as ‘Tagetes oil’, possesses larvicidal properties and is used as a fly repellent. The carotenoids present in *Tagetes* are the major sources of pigments used in poultry industry as a feed additive to intensify the yellow colour of egg yolks and broiler skin (Scott *et al.* 1968). The principal pigment present in the flower is xanthophyll. The xanthophyll, lutein accounts for more than 80–90 per cent and is present in the form of esters of palmitic and myristic acids (Alam *et al.* 1968).

## Materials and Method

The present investigation was carried out as a pot experiment in Department of Agriculture Microbiology, Kittur Rani Channamma College of Horticulture, Arabhavi, Gokak taluk, Belgaum, Karnataka during the period from November 2012–February 2013. The details of the materials used and

the methods adopted during the investigation are presented below.

### Planting Material

The marigold seeds used for the experiment were collected from two sources, a local variety obtained from a farmer in Murgod, Belgaum, and a second variety named Double Orange from Namdhari Seed Company Pvt. Ltd.

### VAM species used for the experiment

- Control
- *Glomus bagyarajii*
- *Glomus fasciculatum*
- *Glomus intraradices*
- *Glomus leptotichum*
- *Glomus monosporum*
- *Glomus monihofis*
- *Glomus mosseae*
- *Entrophospora*
- *Aculospora laevis*
- *Sclerocystis dusii*
- *Gigaspora gigantia*
- *Gigaspora margarita*

The marigold seeds of the two varieties were sown in poly bags with 12 VAM species and a control. Mycorrhiza was placed in the poly bag and then soil was added to it. Marigold seeds were placed over the soil and were finally covered with soil. The poly bags were watered regularly. Weeding was done as per the requirement.

### Inoculation with AM Fungus

Culture of 12 vesicular arbuscular mycorrhizal fungi (VAM) inoculums was obtained from Department of Agricultural Microbiology, Kittur Rani Channamma College of Horticulture, Arabhavi. The inoculum was multiplied in a sterilized potting mixture using maize (*Zea mays*) as host plant in the shade house of Microbiology Department. The inoculum used consisted of sand and soil in 1:1 ratio, and root segments of maize comprising hyphae, vesicles, arbuscules, and chlamydospores of 12 AM fungus. 5 g of inoculum was applied per poly bag before sowing.

\* Department of Agricultural Microbiology, K R C College of Horticulture, Arabhavi 591218. E-mail: patilpradeep@yahoo.co.in



## Results and Discussion

A mutualistic symbiotic association exists between the endomycorrhizal fungi (AM fungi) and the roots of the horticultural crops (host plants). The AM fungal symbiosis starts with the penetration through the radicle and finally to the roots of the germinating seedlings. The local variety of marigold seeds were inoculated with 12 different AM fungal inoculums in the germinating media. The media constituents help in the early germination of the seeds. The germination percentage and vigour was influenced by different VA mycorrhizal fungi on the Local variety of marigold (Figs. 1, 2, 7, and 8). Least number of days taken for germination was recorded by *Gigaspora margarita* (5 days) followed by *Glomus fasciculatum*, *Glomus monihofis*, *Entrophospora*, and *Gigaspora gigantia* (6 days). The highest germination percentage was recorded in *Gigaspora margarita*, *Glomus bagyarajii*, *Glomus monihofis*, *Entrophospora*, and *Gigaspora gigantia* (93.33 per cent).

In Double Orange variety of Marigold, days taken for germination, germination per cent and vigour were also influenced by different VA mycorrhizal fungi existing on it. Least number of days taken for germination was recorded by *Gigaspora margarita*

(5 days) followed by *Sclerocystidusii*, *Entrophospora*, *Glomus monosporum*, *Glomus leptotichum*, and *Glomus fasciculatum* (6 days). The highest germination percentage was recorded in *Gigaspora gigantia* (100 per cent) followed by *Glomus monihofis* (93.33 per cent), when compared to control (Table 1 and 2). The significant enhancement in germination was also noticed in VAM inoculated seeds when compared to uninoculated control (Figs. 3, 4, 5, and 6). Increased germination due to VAM inoculation is also reported in papaya (Duragannavar *et al.* 2004), and in jamun (Devachandra, 2006).

Enhancement in germination could be due to the fact that soon after sowing of seeds they start imbibing water resulting in triphasic increase in seed fresh weight (Hartmann *et al.* 1997). Another important characteristic of seeds during imbibition is that they become 'leaky'. Several compounds including amino acids, organic acids, inorganic ions, sugars, phenolic compounds, and proteins leak out from the imbibing seeds (Simon, 1984). These solutes/leaked compounds may help VAM fungal propagules to germinate early. The time gap between the first phase and the last phase of germination may help the AM fungi to germinate and establish contacts as soon as

Fig 1: Effect of 12 different VAM species on growth of marigold Local variety



Fig 2: Effect of different host preferred VAM on growth of marigold Local variety



Con-Control, G.mono- *Glomus monosporum*, Entro-*Entrophospora*, A.I-*Aculospora laeives*, S.d-*Sclerocystis dussii*

Fig 3: Effect of 12 different VAM species on growth of marigold Double Orange variety



Fig 4: Effect of host preferred VAM on growth of marigold Double Orange variety



Con-Control, Entro-*Entrophospora*, A.I-*Aculospora laeives*, S.d-*Sclerocystis dussii*, Gi.g-*Glomus gigantia*

**Fig 5:** Effect of 12 VAM species on shoot and root growth of marigold Double Orange variety



**Fig 6:** Effect of host preferred VAM on shoot and root growth of marigold Double Orange variety



S.d-Sclerocystis dussii, A.l-Aculospora laeives, Entro-Entrophospora, G.mono Glomus monosporum, G.l-Glomus leptotichum, Con-Control

**Fig 7:** Effect of 12 VAM species on shoot and root growth of marigold Local variety



**Fig 8:** Effect of host preferred VAM on root and shoot growth of marigold Local variety



Con-Control, G.l-Glomus leptotichum, G.moss-Glomus mossaea, Entro Entrophospora, A.l-Aculospora laeives

**Table 1:** Effect of VAM species on days taken for germination and germination percentage in marigold Local variety

Treatments	Days taken for germination	Germination percentage
T <sub>1</sub> - Control	7.00	86.33
T <sub>2</sub> . <i>Glomus bagyarajii</i>	8.00	93.63
T <sub>3</sub> . <i>Glomus fasciculatum</i>	6.00	66.66
T <sub>4</sub> . <i>Glomus intraradices</i>	8.00	86.66
T <sub>5</sub> . <i>Glomus leptotichum</i>	8.00	73.33
T <sub>6</sub> . <i>Glomus monosporum</i>	8.00	66.66
T <sub>7</sub> . <i>Glomus monihofis</i>	6.00	93.33
T <sub>8</sub> . <i>Glomus mosseae</i>	7.00	86.66
T <sub>9</sub> . <i>Entrophospora</i>	6.00	93.33
T <sub>10</sub> . <i>Aculospora laevis</i>	8.00	80.00
T <sub>11</sub> . <i>Sclerocystis dusii</i>	7.00	80.00
T <sub>12</sub> . <i>Gigaspora gigantea</i>	6.00	93.33
T <sub>13</sub> . <i>Gigaspora margarita</i>	5.00	93.33
Standard Error Mean (S Em) ±	0.06	0.21
Critical Difference (C. D) @1 %	0.23	0.82

**Table 2:** Effect of VAM species on days taken for germination and germination per cent in marigold Double Orange variety

Treatments	Days taken for germination	Germination percentage
T <sub>1</sub> - Control	9.00	80.03
T <sub>2</sub> . <i>Glomus bagyarajii</i>	9.00	73.33
T <sub>3</sub> . <i>Glomus fasciculatum</i>	6.00	80.00
T <sub>4</sub> . <i>Glomus intraradices</i>	8.00	86.66
T <sub>5</sub> . <i>Glomus leptotichum</i>	6.00	80.00
T <sub>6</sub> . <i>Glomus monosporum</i>	6.00	73.33
T <sub>7</sub> . <i>Glomus monihofis</i>	8.00	93.33
T <sub>8</sub> . <i>Glomus mosseae</i>	5.00	73.33
T <sub>9</sub> . <i>Entrophospora</i>	6.00	86.66
T <sub>10</sub> . <i>Aculospora laevis</i>	6.00	73.33
T <sub>11</sub> . <i>Sclerocystis dusii</i>	9.00	66.66
T <sub>12</sub> . <i>Gigaspora gigantea</i>	7.00	100.00
T <sub>13</sub> . <i>Gigaspora margarita</i>	5.00	80.00
S Em ±	0.06	0.01
C. D @1 %	0.23	0.04

radicles emerge out. Hence, AM fungi help better seed germination by mutualistic symbiosis with roots of the plant and competing with the pathogens for space and nutrition. The differences observed in the efficacy of germination by the different AM fungal species could be attributed to the leached solutes leaked from the appropriate host. Thus, leachet released might play a prominent role in early propagation of AM fungal mycelium, thereby contributing to improved efficiency and early seed germination (Cruz *et al.* 2003).

These plant exudates enhance the hyphal growth as the fungus approach the vicinity of the roots. However, these promoting effects appear to be host specific.

In the Local variety (Table 3), highest shoot height was recorded in *Glomus mosseae* (41.90 cm) as compared to control (21.63 cm). Highest root length was recorded in *Glomus monosporum* (36.00 cm) as compared to control (15.13 cm). Highest number of roots was recorded in *Glomus mosseae* (66.33) followed by *Acaulospora laevis* (51.33), and least in *Glomus bagyarajii* (18.33). Highest root volume was recorded in *Glomus leptotichum* (3.97 ml) followed by *Glomus monihofis* (3.50 ml), *Glomus mosseae* (2.83 ml), and *Acaulospora laevis* (3.77 ml) as compared to control (0.43 ml). Highest fresh root weight was recorded in *Glomus leptotichum* (5.40 mg) and least in control (0.15 mg). Highest fresh shoot weight was recorded in *Glomus mosseae* (9.44 mg) and least in the control (0.83 mg). Highest root to shoot ratio was recorded

in *Glomus monihofis* (0.81), which was on par with *Glomus leptotichum* (0.63). The least root to shoot weight was recorded in both *Gigaspora margarita* and *Sclerocystis dusii* (0.15).

In the Double Orange variety (Table 4), the highest shoot height was recorded in *Entrophospora* (36.00 cm) on par with all the other treatments. The least shoot height was recorded in control (22.83 cm). Highest root length was recorded in *Glomus fasciculatum* (36.33cm) and the least in control (15.00cm). Highest number of roots was recorded in *Entrophospora* (58.33) which was on par with all the other treatment other than control (22.33). Highest root volume was recorded in *Entrophospora* (5.33 ml) which was on par with *Glomus leptotichum* (4.83 ml), *Acaulospora laevis* (4.33 ml), *Glomus intraradices* (4.17 ml), *Glomus fasciculatum* (3.17 ml), and *Glomus mosseae* (3.50ml). The least root volume was recorded in control (0.67 ml). Highest fresh root weight was recorded in *Entrophospora* (5.78 mg) and least in control (0.51 mg). Highest fresh shoot weight was recorded in *Glomus mosseae* (8.33 mg) and the least in control (3.33 mg). Highest root to shoot ratio was recorded in *Entrophospora* (0.72) and the least in *Gigaspora gigantia* (0.20).

In the present investigation on marigold, the morphological and anatomical changes were evident with respect to shoot parameters, such as height of shoot and number of leaves, as well as to root parameters, such as number of roots, length of longest

**Table 3:** Effect of 12 species of VAM on growth of Marigold Local variety

Treatments	Shoot length (cm)	Root length (cm)	No. of leaves	No. of roots	Root volume (ml)	Fresh root weight (mg)	Fresh shoot weight (mg)	Root: shoot ratio
T <sub>1</sub> - Control	21.63	15.13	10.67	26.33	0.43	0.15	0.83	0.40
T <sub>2</sub> . <i>Glomus bagyarajii</i>	22.00	24.93	8.33	18.33	1.10	0.62	2.68	0.22
T <sub>3</sub> . <i>Glomus fasciculatum</i>	24.63	15.37	10.00	28.33	1.13	0.71	2.93	0.24
T <sub>4</sub> . <i>Glomus intraradices</i>	30.50	22.43	11.33	28.33	2.00	0.97	5.27	0.46
T <sub>5</sub> . <i>Glomus leptotichum</i>	35.67	28.33	11.67	44.33	3.97	5.40	8.12	0.65
T <sub>6</sub> . <i>Glomus monosporum</i>	27.50	36.00	10.00	41.33	2.17	1.82	3.69	0.45
T <sub>7</sub> . <i>Glomus monihofis</i>	31.50	28.33	10.50	33.33	3.50	4.45	5.47	0.81
T <sub>8</sub> . <i>Glomus mosseae</i>	41.90	33.17	12.67	66.33	2.83	4.39	9.44	0.46
T <sub>9</sub> . <i>Entrophospora</i>	36.67	22.00	12.00	30.67	2.50	1.98	7.36	0.27
T <sub>10</sub> . <i>Acaulospora laevis</i>	34.50	30.83	11.67	51.33	3.77	3.60	8.13	0.41
T <sub>11</sub> . <i>Sclerocystis dusii</i>	30.83	18.30	9.67	25.67	1.83	0.78	5.29	0.15
T <sub>12</sub> . <i>Gigaspora gigantia</i>	31.67	15.67	10.33	33.33	1.23	0.97	6.27	0.16
T <sub>13</sub> . <i>Gigaspora margarita</i>	27.00	20.00	11.00	24.00	0.83	0.64	4.25	0.15
S Em ±	2.99	2.91	1.55	5.34	0.31	0.05	0.08	0.06
C. D@1 %	11.76	11.43	NS	20.98	1.21	0.19	0.32	0.23

**Table 4:** Effect of 12 species of VAM on growth of marigold Double Orange variety

Treatments	Shoot length (cm)	Root length (cm)	No. of roots	No. of leaves	Root volume (ml)	Fresh root weight (mg)	Fresh shoot weight (mg)	Root : shoot ratio
T <sub>1</sub> - Control	22.83	15.00	22.33	12.00	0.67	0.51	3.33	0.28
T <sub>2</sub> - <i>Glomus bagyarajii</i>	25.17	20.83	42.67	11.33	1.17	0.89	3.66	0.24
T <sub>3</sub> - <i>Glomus fasciculatum</i>	29.33	36.33	44.67	12.00	3.17	2.76	8.33	0.33
T <sub>4</sub> - <i>Glomus intraradices</i>	30.33	24.83	53.67	12.67	4.17	3.20	7.66	0.42
T <sub>5</sub> - <i>Glomus leptotichum</i>	34.00	33.00	55.33	11.33	4.83	3.65	6.66	0.55
T <sub>6</sub> - <i>Glomus monosporum</i>	29.00	27.17	41.67	10.67	1.83	1.88	5.33	0.35
T <sub>7</sub> - <i>Glomus monihofis</i>	29.17	28.17	50.00	13.67	1.83	1.47	5.97	0.24
T <sub>8</sub> - <i>Glomus mosseae</i>	31.83	32.67	52.67	12.00	3.50	2.88	8.33	0.35
T <sub>9</sub> - <i>Entrophospora</i>	36.00	32.33	58.33	12.00	5.33	5.78	7.97	0.72
T <sub>10</sub> - <i>Aculospora laevis</i>	33.33	35.00	48.67	12.67	4.33	2.91	6.33	0.46
T <sub>11</sub> - <i>Sclerocystis dusii</i>	35.33	29.00	39.33	11.33	1.83	2.00	6.33	0.32
T <sub>12</sub> - <i>Gigaspora gigantia</i>	31.00	21.50	45.33	11.33	1.33	1.14	5.66	0.20
T <sub>13</sub> - <i>Gigaspora margarita</i>	31.33	26.33	39.00	10.67	2.33	1.59	5.33	0.30
S Em ±	2.76	2.90	5.69	1.03	0.59	0.19	0.01	0.03
C. D @1 %	10.84	11.40	22.37	NS	2.33	0.73	0.06	0.14

lateral roots, root volume, fresh root weight, fresh shoot weight, and root to shoot ratio (Table 3 and 4).

Modification in the root geometry and morphology might have a morphogenetic effect mediated by Indole Acetic Acid (IAA) and gibberellins (Allen *et al.* 1980). Hooker and Artkinson (1992) concluded that root morphogenesis can be modified by AM fungal metabolism or by hormones independent of external nutrients supplied. Hence, the root morphogenetic growth (i.e., root geometry) and the promising effect of AM fungi observed in the present investigation and from the literature (Slankis, 1957) could also be attributed to the phytohormones, such as gibberellins and auxins, and vitamins produced by the AM fungi. Further, it is well documented that infection of plant roots by AM fungi has beneficial effects on vegetative parameters and biomass production (fresh weight) of host plants (Duragannavar, 2005).

The increase in these parameters may be attributed to the synthesis of hormones and growth factors by AM fungi, leading to increased cell multiplication and cell division resulting in overall increase in plant height and number of leaves.

Percent root colonization was higher in the entire VAM inoculated seedling as compared to control in both the varieties (Table 5). In Local variety, maximum percentage of root colonization was recorded in *Gigaspora margarita*, *Sclerocystis dusii*,


**Table 5:** Effect of VAM inoculation on root colonization

Treatments	Percent root colonization	
	Local	Double Orange
T <sub>1</sub> - Control	55.00	46.00
T <sub>2</sub> - <i>Glomus bagyarajii</i>	81.00	69.00
T <sub>3</sub> - <i>Glomus fasciculatum</i>	86.00	74.00
T <sub>4</sub> - <i>Glomus intraradices</i>	93.00	78.00
T <sub>5</sub> - <i>Glomus leptotichum</i>	98.00	81.00
T <sub>6</sub> - <i>Glomus monosporum</i>	95.66	85.00
T <sub>7</sub> - <i>Glomus monihofis</i>	99.00	90.00
T <sub>8</sub> - <i>Glomus mosseae</i>	99.00	92.00
T <sub>9</sub> - <i>Entrophospora</i>	97.00	87.00
T <sub>10</sub> - <i>Aculospora laevis</i>	98.00	89.00
T <sub>11</sub> - <i>Sclerocystis dusii</i>	99.00	90.00
T <sub>12</sub> - <i>Gigaspora gigantia</i>	98.00	91.00
T <sub>13</sub> - <i>Gigaspora margarita</i>	99.00	79.00
S Em ±	0.58	0.57
C D @1 %	2.29	2.26

*Glomus mosseae* and *Glomus monihofis* (99.00 per cent) which was at par with *Gigaspora gigantea*, *Aculospora laevis*, *Glomus leptotichum* (98.00 per cent), and *Entrophospora* (97.00 per cent), as compared to the control (55.00 per cent). Whereas in Double Orange variety, the maximum percentage of root colonization was recorded in *Glomus mosseae* (92.00 per cent) which was at par with *Glomus mosseae* (92.00 per cent), *Gigaspora gigantea* (91.00 per cent), and *Glomus monihofis* (90.00 per cent). The least root colonization was recorded in the control (46.00 per cent).


The highest root infection/colonization (Fig. 9) and increased root geometry in *Entrophospora*, *Glomus mosseae*, *Aculospora laevis*, *Glomus leptotichum*, *Aculospora laevis*, and *Sclerocystis dussii* inoculated marigold seedlings have direct effect on germination and in the development of sound and healthy rhizosphere that would contribute to improved growth, resulting in improved nutrient uptake. Thus, inoculated seedlings are most suited to grow marigold. Similar results were also reported by Hooker *et al.* (1992); Venkat, (2004); and Santosh, (2004).

Fig 9: Root staining of marigold inoculated with host preferred VAM Fungi



## References

- Alam A U, Couch J R, and Creger C R. 1968. **Fatty acid composition of the xanthophyll esters of *Tagetes erecta* petals.** *Lipids* 3: 183–184
- Allen, M F, Moore T S, and Christensen M. 1980. **Phytohormone changes in *Bouteloua gracillis* infected by vesicular-arbuscular mycorrhiza cytokinin increase in the host plants.** *Canadian J. Botany* 58: 371–374
- Cruz A F, Ishii T, Matsumotto I I, and Kadoya K. 2003. **Evaluation of the mycelia network formed by arbuscular mycorrhizal hyphae in the rhizosphere of papaya and other plants under intercropping system.** *Brazilian Journal of Microbiology* 34(1): 17–21
- Devachandra N. 2006. **Response of jamun to arbuscular mycorrhizal fungi (AM Fungi) and bioformulations on germination, growth of rootstock, graft-take and tolerance to induced moisture stress.** *M Sc (Hort.) Thesis*, Uni. Agri. Sci., Dharwad
- Duragannavar M P, Patil C P, Patil P B, Swamy G S K, and Sidaramayya M B. 2004. **Combined effect of *Glomus fasciculatum* and GA3 treatment on papaya seed germination.** *Thirteen Southern Regional Conference on Microbial Inoculants*, December 3–5, 2004, College of Agriculture, Bijapur, pp 29
- Duragannavar M P. 2005. **Effect of bioformulations on growth and yield of papaya cv. Red lady.** *M Sc (Hort.) Thesis*, Uni. Agri. Sci, Dharwad
- Hartmann H T, Kester D E, Daview F T Jr, and Creneves R L. 1997. **Principles of propagation by seeds.** In *Plant Propagation—Principles and Practices*. Prentice hall of India Private limited, New Delhi, pp 177–194
- Hayman D S. 1980. **Mycorrhiza and crop production.** *Nature* 287: 487–488
- Hooker J E and Atkinson D. 1992. **Application of computer aided image analysis to studies of arbuscular endomycorrhizal fungi effects on plant root system morphology and dynamics.** *Agronomie* 12: 821–824
- Hooker J E, Munro M, and Atkinson D. 1992. **Vesicular arbuscular mycorrhizal fungi induced alteration in polar root system morphology.** *Plant and Soil* 145: 207–214
- Patil P B and Patil C P. 2008. **Mycorrhizal biotechnology for increasing growth and productivity of fruit plants.** In *The mycorrhizae: Diversity, ecology and applications*. Eds. Tiwari M and Sati S C, pp 57–87, Daya publishing House, New Delhi
- Santosh. 2004. **Enhancement of germination, growth, graft rate and stress tolerance of mango rootstocks using bioformulation.** *MSc (Hort.) Thesis*, Univ. Agri. Sci., Dharwad
- Scott M I, Ascarelli I, and Olson G. 1968. **Studies on egg-yolk pigmentation.** *Poultry Science* 47: 863–872
- Simon E W. 1984. **Early events in germination.** In *Seed Physiology*, Academic Press, New York, 2: 77–115
- Slankis V. 1957. **Hormonal relationship in mycorrhizal development.** In *Ectomycorrhizae*, Eds. Marx G C and Kozlowski T T, Academic Press, New York, pp 231–298
- Venkat. 2004. **Exploitation of Rangpur lime as a root stock for different *Citrus* sp.** *M Sc (Hort.) Thesis*, Univ. Agri. Sci., Dharwad



## CENTRE FOR MYCORRHIZAL CULTURE COLLECTION

### Morphotaxonomy of Arbuscular Mycorrhizal Fungi *Acaulospora foveata*

Chaitali Bhattacharya\* and Alok Adholeya#

From this edition we shall commence a new series wherein we would be describing each of our precious Arbuscular Mycorrhizal Fungi cultures available in the Centre for Mycorrhizal Culture Collection Bank on a period basis. According to the present classification studies using ribosomal-RNA (rRNA) gene as marker, phylum Glomeromycota has four orders, viz., Archaeosporales, Diversisporales, Glomerales, and Paraglomerales, comprising 10 families and 13 genera. In this periodical, we have given detailed information about *Acaulospora foveata*, a species with undermentioned classification status (<http://www.zor.zut.edu.pl/Glomeromycota/Classification.html>).

**Glomeromycota** C. Walker & Schuessler

**Glomeromycetes** Cavalier-Smith

**Diversisporales** C. Walker & Schuessler


**Acaulosporaceae** J.B. Morton & Benny

***Acaulospora*** Gerd. & Trappe emend. S.M. Berch

***foveata*** Janos and Trappe

Establishing monosporal (single species) culture for *Acaulospora* spp. was a very challenging process. It took almost two years to get the desired sporulation count (around 50 spores per gram soil). These spores were then isolated for the morphotaxonomic studies (Fig. 1). Undermentioned are the morphotaxonomic features of the organism visualized through voucher specimens study, presented in a similar pattern as adopted by the listed references and sites provided at the end of the article.


Fig 1.: Spores isolated from monosporal culture of *Acaulospora foveata*



#### Spore Morphology

The spores of *Acaulospora foveata* occur singly in the soil (Fig. 2). As reported by many authors, the ontogenesis of the spore begins with the development of a sporiferous saccule at the top of a sporogenous hypha, which is in continuation with the extraradical hyphae. After the full growth of saccule, the spore begins to develop from the side of the subtending hypha. Spore growth can be observed only when the sample is captured at the right stage under suitable *in-situ* conditions (Fig. 3). Once the spore matures, the saccule loses its contents and eventually slashes off from the fully matured spores.

Fig. 2: Singly occurring spores



\* Fellow, Biotechnology and Bioresources, Centre for Mycorrhizal Research, The Energy and Resources Institute, Darbari Seth Block, IHC Complex, Lodhi Road, New Delhi-110 003, India

# Director, Biotechnology and Bioresources, Centre for Mycorrhizal Research, The Energy and Resources Institute, Darbari Seth Block, IHC Complex, Lodhi Road, New Delhi-110 003, India


Fig. 3: Sporiferous saccule



### Spore Colour

The newly formed immature spores are initially cream coloured. On reaching to maturity, the spores acquire a dark red and finally a dark brown colour (Fig. 4).

Fig. 4: Single spores representing size



### Spore Shape

Spores are acaulosporoid. Spore shape is mostly globose to subglobose. The spore size is more than 150 µm. Size distribution ranges between 210–378 µm.

#### Description of Subcellular Structure of Spore


The matured spore wall is composed of three layers (L1, L2, and L3) (Fig. 5). Błaszowski (2012) has reported the undermentioned details of the spore layer in his book chapter.

**L1:** This layer forms the surface of the spore. It is evanescent and hyaline and remains attached to the layer 2. It is completely sloughed in mature spores. The thickness of the wall ranges between 2–3.63 µm.

**L2:** The second layer helps in providing structural integrity to the spore due to the presence of laminae.

A mature second layer has a thickness of 9–19 µm. It is ornamented with evenly distributed circular pits

Fig. 5: Three different wall layers of spore



when observed in plain view (Fig. 6).

**L3:** The third layer is semi-flexible. It is 3–5 µm thick and is usually tightly adhered to the lower surface of layer 2.

### Germinal Walls

There are two hyaline flexible inner walls (gw1 and gw2).

#### Reaction with Melzer's Reagent

It stains pinkish red to a reddish-purple or pastel red with Melzer's reagent (Fig. 7).

Fig. 6: Pitted surface on the spore wall





Fig. 7: Pinkish red stain with Melzer's reagent



### Sporiferous Saccule

It consists of a saccule and a neck. It is normally hyaline, light yellow to apricot yellow in colour, globose to subglobose shaped, has a diameter of 200–250 µm, with neck tapering from the saccule towards its end. It usually collapses at maturity and becomes detached from the mature spores.

Fig. 8: Spore showing cicatrix




### Cicatrix

It is circular to ovoid scar, indicating region of contact between spore and saccule neck. It has a slightly raised tapering collar consisting of closely packed tubercles surrounding an unornamented depression. It is about 12–16 µm in diameter (mean = 13.8 µm) (Fig. 8).

Since morphotaxonomy is not considered to be an authentic tool for identification, we are in the process of undertaking molecular as well as biochemical characterization of the spores isolated from the monosporal cultures for establishing its authenticity.

### References

Janos D P and Trappe J M. 1982. **Two new *Acaulospora* species from tropical America.** *Mycotaxon* 15: 515–522  
Błaszowski J (ed). 2012. *Glomeromycota*, W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków , pp 303  
<http://invam.wvu.edu/the-fungi/classification/acaulosporaceae/acaulospora/foveata>  
<http://www.zor.zut.edu.pl/Glomeromycota/Classification.html>



# Recombinant DNA Technology

Keya Chaudhuri

2013 • ISBN: 9788179933206  
 Pages: 344 • Binding: Paperback  
 Size: 180 × 240 mm • Price: ₹595.00

*Recombinant DNA Technology* is focussed on the current state of knowledge on the recombinant DNA technology and its applications. The book will provide comprehensive knowledge on the principles and concepts of recombinant DNA technology or genetic engineering, protein expression of cloned genes, PCR amplification of DNA, RFLP, AFLP and DNA fingerprinting and finally the most recent siRNA technology. It can be used by post-graduate students studying and teachers teaching in the area of Molecular Biology, Biotechnology, Genetics, Microbiology, Life Science, Pharmacy, Agriculture and Basic Medical Sciences.

## Key features

- Comprehensive analysis of the DNA structure and the process of DNA replication
- Extensive analysis of restriction endonucleases, sequencing by Sanger's method, and protein production in bacteria
- Detailed description of PCR, RFLP, AFLP, and DNA fingerprinting
- Thorough explanation of site-directed mutagenesis and cloning in cosmid vectors
- In-depth discussion on methods for creating recombinant DNA molecules and construction of DNA libraries
- Concepts and applications of agarose gel and polyacrylamide gel electrophoresis

## Table of contents

Recombinant DNA Technology • Methods for Creating Recombinant DNA Molecules • Properties of Restriction Endonucleases • Screening of Recombinant DNA Molecules • Construction of DNA Library • Sequencing by Sanger's Method • Protein Production in Bacteria • Site-directed Mutagenesis • Restriction Fragment Length Polymorphism • Polymerase Chain Reaction DNA Fingerprinting • RNAi and siRNA Technology • Molecular Biology Methods • Features of Commonly Used Vectors • Isolation and Purification of Plasmid Vectors • Cloning in Cosmid Vectors • Construction of Genomic DNA Libraries in Cosmid Vectors • Enzymes Used in Molecular Cloning • Agarose Gel and Polyacrylamide Gel Electrophoresis • Detection and Extraction of DNA from Gels

The Energy and Resources Institute  
 Attn: TERI Press  
 Darbari Seth Block  
 IHC Complex, Lodhi Road  
 New Delhi – 110 003/India

Tel. 2468 2100 or 4150 4900  
 Fax: 2468 2144 or 2468 2145  
 India +91 • Delhi (0)11  
 Email: teripress@teri.res.in  
 Web: <http://bookstore.teriin.org>

To purchase the book, visit our online bookstore at <http://bookstore.teriin.org> or send us your demand draft or cheque in favour of TERI, payable at New Delhi (outstation cheques are not accepted).



## RECENT REFERENCES

The latest additions to the network's database on mycorrhiza are published here for the members' information. The list consists of papers from the following journals:

- *Applied Soil Ecology*
- *Biological Conservation*
- *Crop Protection*
- *Environmental and Experimental Botany*
- *Forest Ecology and Management*
- *Journal of Plant Physiology*
- *Journal of Theoretical Biology*
- *Microbiological Research*
- *Mycorrhiza*
- *Mycoscience*
- *Phytochemistry*
- *Plant Physiology and Biochemistry*
- *Scientia Horticulturae*
- *Soil and Tillage Research*
- *Soil Biology and Biochemistry*
- *South African Journal of Botany*

Name of the author(s) and year of publication	Title of the article, name of the journal, volume number, issue number, page numbers (address of the first author or of the corresponding author, marked with an asterisk)
Adeleke R*, Dames J F. 2014	<b><i>Kalaharituber pfeilii</i> and associated bacterial interactions</b> <i>South African Journal of Botany</i> <b>90</b> : 68–73 [ARC-Institute for Soil, Climate & Water, 600, Belvedere Street, Arcadia 0001, Pretoria]
Calonne M*, Fontaine J, Debiane D, Laruelle F, Grandmougin-Ferjani A, Sahraoui A L-H. 2014	<b>The arbuscular mycorrhizal <i>Rhizophagus irregularis</i> activates storage lipid biosynthesis to cope with the benzo[a]pyrene oxidative stress</b> <i>Phytochemistry</i> <b>97</b> : 30–37 [Univ Lille Nord de France, F-59000 Lille, France]
Caravaca F*, Ruess L. 2014	<b>Arbuscular mycorrhizal fungi and their associated microbial community modulated by Collembola grazers in host plant free substrate</b> <i>Soil Biology and Biochemistry</i> <b>69</b> : 25–33 [Institute of Biology, Ecology Group, Humboldt-Universität zu Berlin, Philipstrasse 13, 10115 Berlin, Germany]
Chen Y-L*, Zhanga X, Ye J-S, Han H-Y, Wan S-Q, Chen B-D. 2014	<b>Six-year fertilization modifies the biodiversity of arbuscular mycorrhizal fungi in a temperate steppe in Inner Mongolia</b> <i>Soil Biology and Biochemistry</i> <b>69</b> : 371–381 [State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China]
Corredor A H*, Rees K V, Vujanovic V. 2014	<b>Host genotype and health status influence on the composition of the arbuscular mycorrhizal fungi in <i>Salix</i> bioenergy plantations</b> <i>Forest Ecology and Management</i> <b>314</b> : 112–119 [Department of Food and Bioproduct Science, University of Saskatchewan, College of Agriculture and Bioresources, 51 Campus Drive, Saskatoon, SK S7N 5A8, Canada]
Fichtner A*, Oheimb G V, Härdtle W, Wilken C, Gutknecht J L M. 2014	<b>Effects of anthropogenic disturbances on soil microbial communities in oak forests persist for more than 100 years</b> <i>Soil Biology and Biochemistry</i> <b>70</b> : 79–87 [Institute for Natural Resource Conservation, University of Kiel, Olshausenstr. 75, 24118 Kiel, Germany]
Freitas R D O*, Buscardo E, Nagy L, Maciel A B D S, Carrenho R, Luizão R C C. 2014	<b>Arbuscular mycorrhizal fungal communities along a pedo-hydrological gradient in a Central Amazonian terra firme forest</b> <i>Mycorrhiza</i> <b>24</b> (1): 21–32 [Instituto Nacional de Pesquisas da Amazônia (INPA), Coordenação de Biodiversidade (CBIO), Av. Efigênio Sales, 2239, Campus III CP 478, CEP 69011-970, Manaus, Amazonas, Brazil]

Name of the author(s) and year of publication	Title of the article, name of the journal, volume number, issue number, page numbers (address of the first author or of the corresponding author, marked with an asterisk)
Füzy A*, Bothe H, Molnár E, Biró B. 2014	<p><b>Mycorrhizal symbiosis effects on growth of chalk false-brome (<i>Brachypodium pinnatum</i>) are dependent on the environmental light regime</b>  <i>Journal of Plant Physiology</i> 171(5): 1–6            [Institute for Soil Sciences and Agricultural Chemistry, Centre for Agricultural Research, Hungarian Academy of Sciences, Herman Ottó út 15, H-1022 Budapest, Hungary]</p>
García I*, Mendoza R. 2014	<p><b><i>Lotus tenuis</i> seedlings subjected to drought or waterlogging in a saline sodic soil</b>  <i>Environmental and Experimental Botany</i> 98: 47–55            [Museo Argentino de Ciencias Naturales Bernardino Rivadavia-Conicet, Av. Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina]</p>
Abd-Alla M H*, El-Enany AW E, Nafady N A, Khalaf D M, Morsy F M. 2014	<p><b>Synergistic interaction of <i>Rhizobium leguminosarum</i> bv. <i>viciae</i> and arbuscular mycorrhizal fungi as a plant growth promoting biofertilizers for faba bean (<i>Vicia faba</i> L.) in alkaline soil</b>  <i>Microbiological Research</i> 169(1): 49–58            [Department of Botany and Microbiology, Faculty of Science, Assiut University, Assiut, Egypt]</p>
Jones M D*, Nara K, Hamel C. 2014	<p><b>Highlights of the Seventh International Conference on Mycorrhiza in New Delhi—A conference to remember</b>  <i>Mycorrhiza</i> 24(1): 77–78            [International Mycorrhiza Society, Biology Department, University of British Columbia (Okanagan Campus), Sci 385, 3333 University Way, Kelowna, BC, V1V 1V7, Canada]</p>
Koron D*, Sonjak S, Regvar M. 2014	<p><b>Effects of non-chemical soil fumigant treatments on root colonization with arbuscular mycorrhizal fungi and strawberry fruit production</b>  <i>Crop Protection</i> 55: 35–41            [Agricultural Institute of Slovenia, Department of Fruit and Vine Growing, Hacquetova ulica 17, SI-1000 Ljubljana, Slovenia]</p>
Longo S*, Nouhra E, Goto B T, Berbara R L, Urcelay C. 2014	<p><b>Effects of fire on arbuscular mycorrhizal fungi in the Mountain Chaco forest</b>  <i>Forest Ecology and Management</i> 315: 86–94            [Instituto Multidisciplinario de Biología Vegetal (IMBIV), CONICET, Universidad Nacional de Córdoba, Casilla de correo 495, 5000 Córdoba, Argentina]</p>
Mechri B*, Manga A G B, Tekaya M, Attia F, Cheheb H, Meriem F B, Braham M, Boujnah D, Hammami M. 2014	<p><b>Changes in microbial communities and carbohydrate profiles induced by the mycorrhizal fungus (<i>Glomus intraradices</i>) in rhizosphere of olive trees (<i>Olea europaea</i> L.)</b>  <i>Applied Soil Ecology</i> 75: 124–133            [Laboratoire de Biochimie, USCR Spectrométrie de Masse, LR-NAFS/LR12ES05 Nutrition-aliments fonctionnels et santé vasculaire, Faculté de Médecine, Université de Monastir, 5019 Monastir, Tunisia]</p>
Navarro J M*, Pérez-Tornero O, Morte A. 2014	<p><b>Alleviation of salt stress in citrus seedlings inoculated with arbuscular mycorrhizal fungi depends on the rootstock salt tolerance</b>  <i>Journal of Plant Physiology</i> 171(1): 76–85            [Departamento Citricultura, Instituto Murciano de Investigación y Desarrollo Agrario y Alimentario (IMIDA), C/ Mayor s/n, 30150 La Alberca, Murcia, Spain]</p>
Ortas I*, Ustuner O. 2014	<p><b>Determination of different growth media and various mycorrhizae species on citrus growth and nutrient uptake</b>  <i>Scientia Horticulturae</i> 166: 84–90            [Department of Soil Science and Plant Nutrition, University of Çukurova, Faculty of Agriculture, Adana, Turkey]</p>

Name of the author(s) and year of publication	Title of the article, name of the journal, volume number, issue number, page numbers (address of the first author or of the corresponding author, marked with an asterisk)
Pérez-Tienda J*, Corrêa A, Azcón-Aguilar C, Ferrol N. 2014	<b>Transcriptional regulation of host view the NH<sub>4</sub>(+) source transporters and GS/GOGAT pathway in arbuscular mycorrhizal rice roots</b> <i>Plant Physiology and Biochemistry</i> 75: 1–8 [Departamento de Microbiología del Suelo y Sistemas Simbióticos, Estación Experimental del Zaidín, CSIC, C. Profesor Albareda 1, 18008 Granada, Spain]
Phillips R D*, Peakall R, Hutchinson M F, Linde C C, Xu T, Dixon K W, Hopper S D. 2014	<b>Specialized ecological interactions and plant species rarity: The role of pollinators and mycorrhizal fungi across multiple spatial scales</b> <i>Biological Conservation</i> 169: 285–295 [Evolution, Ecology and Genetics, Research School of Biology, The Australian National University, Canberra ACT 0200, Australia]
Talaata N B*, Shawky B T. 2014	<b>Protective effects of arbuscular mycorrhizal fungi on wheat (<i>Triticum aestivum</i> L.) plants exposed to salinity</b> <i>Environmental and Experimental Botany</i> 98: 20–31 [Department of Plant Physiology, Faculty of Agriculture, Cairo University, Giza, Egypt]
Taylor A*, Walker C, Bending G D. 2014	<b>Dimorphic spore production in the genus <i>Acaulospora</i></b> <i>Mycoscience</i> 55(1): 1–4 [Warwick Crop Centre, School of Life Sciences, University of Warwick, Wellesbourne, Warwick CV35 9EF, UK]
Thornley J H M*, Parsons A J. 2014	<b>Allocation of new growth between shoot, root and mycorrhiza in relation to carbon, nitrogen and phosphate supply: Teleonomy with maximum growth rate</b> <i>Journal of Theoretical Biology</i> 342: 1–14 [Centre for Nutrition Modelling, Department of Animal & Poultry Science, University of Guelph, Guelph, Ont., Canada N1G 2W1]
Xie X*, Weng B, Cai B, Dong Y, Yan C. 2014	<b>Effects of arbuscular mycorrhizal inoculation and phosphorus supply on the growth and nutrient uptake of <i>Kandelia obovata</i> (Sheue, Liu &amp; Yong) seedlings in autoclaved soil</b> <i>Applied Soil Ecology</i> 75: 162–171 [Key Laboratory of the Ministry of Education for Coastal and Wetland Ecosystems, Xiamen University, Xiamen 361005, PR China]
Yamada A*, Endo N, Murata H, Ohta A, Fukuda M. 2014	<b><i>Tricholoma matsutake</i> Y1 strain associated with <i>Pinus densiflora</i> shows a gradient of in vitro ectomycorrhizal specificity with Pinaceae and oak hosts</b> <i>Mycoscience</i> 55: 27–34 [Department of Bioscience and Biotechnology, Faculty of Agriculture, Shinshu University, 8304 Minami-minowa, Nagano 399-4598, Japan]

# FORTHCOMING EVENTS

## CONFERENCES, CONGRESSES, SEMINARS, SYMPOSIUMS, AND WORKSHOPS

Napa Valley, USA  
17-20 January, 2014

### Truffle Festival

The Westin Verasa Napa, 1314 McKinstry Street, Napa, CA, USA 94559  
Tel. (888) 753-9378  
E-mail [info@napatrufflefestival.com](mailto:info@napatrufflefestival.com)

Taipei, Taiwan  
4-8 May, 2014

### APMBC 2014

APMBC 2014 Congress Secretariat, c/o K&A International Co., Ltd, 3F., No. 183, Kangchien Rd, Taipei, Taiwan 11494

Tel. +886-2-8751-3588  
Fax +886-2-8751-2799  
E-mail [apmbc2014@knaintl.com.tw](mailto:apmbc2014@knaintl.com.tw)  
Web site <http://www.apmbc2014.com/>

Zurich, Switzerland  
14-16 May, 2014

### 33rd New Phytologist Symposium: Networks of Power and Influence: Ecology and Evolution of Symbioses between Plants and Mycorrhizal Fungi

New Phytologist Central Office, Bailrigg House, Lancaster University, LA1 4YE, UK

Tel. +44 1524 594 691  
Fax +44 1524 594 696  
E-mail [np-symposia@lancaster.ac.uk](mailto:np-symposia@lancaster.ac.uk)  
Web site <http://www.newphytologist.org/registrations/index/symposium/4>

Elenite Holiday Village,  
Burgas, Bulgaria  
5-9 June, 2014

### Agriculture and Food 2014, 2nd International Conference

Ivan Genov  
E-mail [agriculture@sciencebg.net](mailto:agriculture@sciencebg.net)  
Website <http://www.sciencebg.net/en/conferences/agriculture-and-food/>

The Convention Centre  
Dublin, Ireland  
22-26 June, 2014

### Plant Biology Europe FESPB/EPSO Congress 2014

FESPB 2014 Conference Secretariat, c/o Keynote PCO, Suite 26, Anglesea House, 63 Carysfort Ave, Blackrock, Co. Dublin, Ireland

Tel. + 353 (0) 1 400 3626  
Fax + 353 (0) 1 400 3692  
E-mail [registration@europlantbiology.org](mailto:registration@europlantbiology.org)  
Web site <http://europlantbiology.org/>

Bangkok, Thailand  
3-8 August, 2014

### The 10th International Mycological Congress

Ms Annawan Knight, NCC Management and Development Company Limited, 60 Queen Sirikit National Convention Centre, New Rachadapisek Road, Klongtoey Bangkok, 10110 Thailand

Tel. (66) 2 229 3335  
Fax (66) 2 229 3346  
E-mail [IMC10secretariat@gmail.com](mailto:IMC10secretariat@gmail.com), [IMC10bkk@biotec.or.th](mailto:IMC10bkk@biotec.or.th)  
Website <http://www.IMC10.com>

Melbourne, Australia  
10-15 August, 2014

### International Association of Plant Biotechnology Congress 2014

119 Buckhurst Street, South Melbourne VIC 3205 Australia

Tel. +61 3 9645 6311  
Fax +61 3 9645 6322  
E-mail [iapb@wsm.com.au](mailto:iapb@wsm.com.au)  
Web site <http://www.iapb2014congress.com/>

---

Editor Alok Adholeya • Associate Editor T P Sankar • Assistant Editor Hemambika Varma

---